

# *Parity Violation From Few Nucleon Systems*

*Michael Gericke*

*Chiral Dynamics Workshop*

*Bern, July 6-10 2009*

**physics &  
astronomy**

**university of manitoba**



# *Outline*

## *PV and the hadronic weak interaction*

Shi-Lin Zhu, et al. , Nuclear Physics A 748 (2005) 435-498

M.J. Ramsey Musolf and S. Page, Annu. Rev. Nucl. Part. Sci. 2006. 56:1-52

C.-P. Liu, Nuclear Physics Phys. Rev. C 75, 065501 (2007)

## *Meson Exchange Picture*

## *EFT and Hadronic PV*

## *The experimental program*

*Parity violating processes between nucleons are used as a tool to study the hadronic weak interaction (HWI) as well as how it is modified by the strong interactions from the simple Standard Model prediction.*

*Two (common) ways to study HWI:*

*1. Flavor changing  $\Delta S=1$  hyperon and meson decay*

*➤ Decay amplitudes, asymmetries, ...*

*2. Flavor conserving  $\Delta S=0$  PV interactions at low energy*

*➤ Mostly asymmetries, analyzing power, rotation angles*

## *Flavor changing decay of mesons and hyperons:*

- *Much theoretical progress from EFT,  $\chi$ PT, heavy quark EFT*
- *Structure of operators from effective Lagrangians incorporate the symmetries of QCD*

*Not so, in hyperon decay:*

- *Unresolved  $\Delta I = \frac{1}{2}$  rule puzzle*
- *Anomalously large PV asymmetries in hyperon radiative decays*
- *Etc.*

*Do the unexpected observations in the  $\Delta S=1$  sector come from a dynamical strange quark or some other process ?*

 *Look at the  $\Delta S=0$  sector*

**Standard Model:** 
$$\mathcal{L}_W^{INT} = -\frac{g}{2\sqrt{2}} \left( J_C^{\mu\dagger} W_\mu + J_c^\mu W_\mu^\dagger \right) - \frac{g}{4 \cos \theta_w} J_N^\mu Z_\mu$$

**Charged currents:**

$$J_C^\mu = \bar{\psi}_d \gamma^\mu (1 - \gamma_5) \psi_u \cos \theta_c + \bar{\psi}_s \gamma^\mu (1 - \gamma_5) \psi_u \sin \theta_c \\ - \bar{\psi}_d \gamma^\mu (1 - \gamma_5) \psi_c \sin \theta_c + \bar{\psi}_s \gamma^\mu (1 - \gamma_5) \psi_c \cos \theta_c$$

**Neutral currents:**

$$J_N^\mu = \bar{\psi}_u \gamma^\mu \left( 1 - \frac{8}{3} \sin^2 \theta_w - \gamma_5 \right) \psi_u + \bar{\psi}_c \gamma^\mu \left( 1 - \frac{8}{3} \sin^2 \theta_w - \gamma_5 \right) \psi_c \\ - \bar{\psi}_d \gamma^\mu \left( 1 - \frac{4}{3} \sin^2 \theta_w - \gamma_5 \right) \psi_d - \bar{\psi}_s \gamma^\mu \left( 1 - \frac{4}{3} \sin^2 \theta_w - \gamma_5 \right) \psi_s$$

**Do the unexpected observations in the  $\Delta S=1$  sector come from a dynamical strange quark or some other process ?**


**→ Look at the  $\Delta S=0$  sector**

**Standard Model:** 
$$\mathcal{L}_W^{INT} = -\frac{g}{2\sqrt{2}} (J_C^{\mu\dagger} W_\mu + J_c^\mu W_\mu^\dagger) - \frac{g}{4 \cos \theta_w} J_N^\mu Z_\mu$$

**Charged currents:**

$$J_C^\mu = \bar{\psi}_d \gamma^\mu (1 - \gamma_5) \psi_u \cos \theta_c + \bar{\psi}_s \gamma^\mu (1 - \gamma_5) \psi_u \sin \theta_c - \bar{\psi}_d \gamma^\mu (1 - \gamma_5) \psi_c \sin \theta_c + \bar{\psi}_s \gamma^\mu (1 - \gamma_5) \psi_c \cos \theta_c$$

*beta decay =>*



$\Delta S = \pm 1$

**Neutral currents:**

$$J_N^\mu = \bar{\psi}_u \gamma^\mu (1 - \frac{8}{3} \sin^2 \theta_w - \gamma_5) \psi_u + \bar{\psi}_c \gamma^\mu (1 - \frac{8}{3} \sin^2 \theta_w - \gamma_5) \psi_c - \bar{\psi}_d \gamma^\mu (1 - \frac{4}{3} \sin^2 \theta_w - \gamma_5) \psi_d - \bar{\psi}_s \gamma^\mu (1 - \frac{4}{3} \sin^2 \theta_w - \gamma_5) \psi_s$$

$\Delta S = 0$   
Hadronic weak interaction!

## *Goals of $\Delta S=0$ HWI studies:*

- 1. Answer how the symmetries of QCD characterize the HWI in strongly interacting systems*

*The HWI is just a residual effect of the  $q$ - $q$  weak interaction for which the range is set by the mass of the  $Z, W$  bosons which is much smaller than the size of nucleons, as determined by QCD dynamics*

 *HWI probes short range  $qq$  correlations*

- 2. Shed light on the puzzles in the  $\Delta S=1$  sector of the HWI*

$Q_{Weak}^p$  measures the electron beam helicity correlated asymmetry in the number of elastically scattered electrons from protons in a liquid hydrogen target at very forward angles, to extract the weak charge of the proton.

$$A_{LR}(\vec{e}, p) = \frac{d\sigma_L - d\sigma_R}{d\sigma_L + d\sigma_R} = k(A_{Q_W^p} + A_{H,V} + A_{H,A})$$


$$A_{Q_W^p} = Q^2 Q_W^p$$

Quantity of interest = -0.288 ppm

$$A_{H,V} = Q_W^n \frac{\epsilon G_E^{p,\gamma} G_E^{n,\gamma} + \tau G_M^{p,\gamma} G_M^{n,\gamma}}{\epsilon (G_E^{p,\gamma})^2 + \tau (G_M^{p,\gamma})^2} + Q_W^s \frac{\epsilon G_E^{p,\gamma} G_E^s + \tau G_M^{p,\gamma} G_M^s}{\epsilon (G_E^{p,\gamma})^2 + \tau (G_M^{p,\gamma})^2}$$

$$A_{H,A} = Q_W^e \frac{\epsilon' G_A^{p,Z} G_M^{p,\gamma}}{\epsilon (G_E^{p,\gamma})^2 + \tau (G_M^{p,\gamma})^2}$$

Must know this from world data:



Axial form factor due to q-q weak interaction related to NN experimental results

Hadronic structure: Must know hadronic wave function or measured form-factors



## $\Delta S=0$ HWI studies:

The  $\Delta S=0$  HWI can only be isolated experimentally via PV observables, to isolate the weak interaction from the much larger EM and strong interactions.

$$\frac{g_W^2}{\alpha M_W^2} \approx 10^{-4}$$

*Weak e-N scale*

$$\frac{g_W^2}{M_W^2} \cdot \frac{M_\pi^2}{g_{\pi NN}^2} \approx 10^{-7}$$

*Weak N-N scale*

*Very challenging !  NIMP experiments*

## $\Delta S=0$ HWI studies:

The  $\Delta S=0$  HWI can only be isolated experimentally via PV observables, to isolate the weak interaction from the much larger EM and strong interactions.

$$\frac{g_W^2}{\alpha M_W^2} \approx 10^{-4}$$

*Weak e-N scale*

$$\frac{g_W^2}{M_W^2} \cdot \frac{M_\pi^2}{g_{\pi NN}^2} \approx 10^{-7}$$

*Weak N-N scale*

*Very challenging !*



*NIMP experiments*



*Nearly Impossible*

## $\Delta S=0$ HWI studies:

*So people started to look for nuclear many-body (large  $A$ ) systems for which there exists some fortuitous enhancement of the size of the observable:*



*coming from nearly degenerate opposite parity state mixing and interference with the much larger parity allowed transition in nuclear excited states.*

*e.g. TRIPLE collaboration:*


*parity violation in compound nuclei from neutron-nucleus resonant scattering with longitudinal cross section asymmetries of order  $10^{-3}$ - $10^{-1}$  (up to  $10^6$  enhancement)*

*But you can get the weak spreading width (weak mixing amplitude) from statistical analysis of this data:*

➔
 $\Gamma_W = 1.8_{-0.3}^{+0.4} \times 10^{-7} \text{ eV}$ 
 $\left( \frac{\langle \psi_f | W | \psi_i \rangle}{\Delta E} \approx \sqrt{\frac{\Gamma_W}{2\pi D}} \right)$

*Can also have large(r) asymmetries from neutron radiative capture (here Cl):* M.T. Gericke *et al.* Phys. Rev. C 74, 065503 (2006)

$$A_\gamma = (-19 \pm 2) \times 10^{-6} \text{ and } A_{LR} = (-1 \pm 2) \times 10^{-6}$$



*n-capture  
on Chlorine  
( $CCl_4$ )*

## $\Delta S=0$ HWI studies:

*However, many-body systems are hard to deal with when it comes to interpretation of the results in a non-statistical fashion.*


*There is no transparent connection to SM.*

 *So back to few body systems*

- No nuclear structure physics*
- Low nucleon momentum ( $\leq \sim 40$  MeV) allows for EFT momentum expansion*
- But no enhancement of asymmetries*

 *Need better experiments*

# The Nucleon-Nucleon Weak Interaction Meson Exchange "Traditional" Picture



$$H_{PC} = ig_{\pi NN} \int d^3x \bar{\psi}_i(x) \gamma^5 \psi_j(x) (\vec{\tau} \cdot \vec{\phi}(x))$$

$$H_{PNC} = \frac{h_\pi^1}{\sqrt{2}} \int d^3x' \bar{\psi}_i(x') \psi_j(x') (\vec{\tau} \times \vec{\phi}(x'))$$

Solutions to the Lippmann-Schwinger equation -  
Essentially the first order term in a Born series:

$$\langle f | V_{PNC} | i \rangle = \langle N_f N_f | H_{PC} \frac{1}{E_0 - H_0 + i\epsilon} H_{PNC} | N_i N_i \rangle$$



$$\frac{ig_{\pi NN} h_\pi^1}{\sqrt{32M}} [\vec{\tau}_1 \times \vec{\tau}_2]_z [\vec{\sigma}_1 + \vec{\sigma}_2] \cdot \left[ \vec{p}, \frac{e^{-mr}}{4\pi r} \right]$$

Weak  $\pi$ -Nucleon Coupling  
( $\rho, \omega$  not shown)

## *Meson exchange picture cont.*

$$\begin{aligned} & \langle N_f N_f | H_{PC} \frac{1}{E_0 - H_0 + i\varepsilon} H_{PNC} | N_i N_i \rangle \\ &= \sum_I \int \frac{d^3 k}{(2\pi)^3} \langle N_f | S | N_i, \pi_I(k) \rangle \frac{1}{\omega_k} \langle N_f, \pi_I(k) | S | N_i \rangle \end{aligned}$$

## Meson exchange picture cont.

$$\begin{aligned} & \langle N_f N_f | H_{PC} \frac{1}{E_0 - H_0 + i\varepsilon} H_{PNC} | N_i N_i \rangle \\ &= \sum_I \int \frac{d^3 k}{(2\pi)^3} \langle N_f | S | N_i, \pi_I(k) \rangle \frac{1}{\omega_k} \langle N_f, \pi_I(k) | S | N_i \rangle \end{aligned}$$

Relationship to quark degrees of freedom:

$$S = \frac{i}{2} \int dt_1 \int dt_2 \int dt_3 \{ H_W^I(t_1) H_W^I(t_2) H_S^I(t_3) \}$$

$$H_W^I = \int d^3 x \left[ \frac{g}{2\sqrt{2}} (J_C^{\mu*} W_\mu + J_C^\mu W_\mu^*) + \frac{g}{4 \cos \theta_w} J_N^\mu Z_\mu \right]$$

$$H_S^I = - \int d^3 x \int d^3 y \int dt_y [ J_S f(x, y) \delta(t_x - t_y) ]$$



## Meson exchange picture cont.

$$\langle N_f N_f | H_{PC} \frac{1}{E_0 - H_0 + i\varepsilon} H_{PNC} | N_i N_i \rangle$$

$$= \sum_I \int \frac{d^3 k}{(2\pi)^3} \langle N_f | S | N_i, \pi_I(k) \rangle \frac{1}{\omega_k} \langle N_f, \pi_I(k) | S | N_i \rangle$$

Relationship to quark degrees of freedom:

~~$$S = \frac{i}{2} \int dt_1 \int dt_2 \int dt_3 \{ H_W^I(t_1) H_W^I(t_2) H_S^I(t_3) \}$$~~

~~$$H_W^I = \int d^3 x \left[ \frac{g}{2\sqrt{2}} (J_C^{\mu*} W_\mu + J_C^\mu W_\mu^*) + \frac{g}{4 \cos \theta_w} J_N^\mu Z_\mu \right]$$~~

~~$$H_S^I = - \int d^3 x \int d^3 y \int dt_y [ J_S f(x, y) \delta(t_x - t_y) ]$$~~

*DDH use SU(6), quark model, and measured hyperon decay amplitudes instead!*

# DDH Model - Benchmark

B. Desplanques, J.F. Donoghue, B.R. Holstein, *Annals of Physics* 124:449-495 (1980)

Arrive at 7 weak meson-nucleon couplings:

$h_{\pi}^1, h_{\rho}^{0,1,2}, h_{\omega}^{0,2}, h_{\rho}^{1'}$

calculated later by Holstein  $\approx 1.8 g_{\pi}$

PV coupling	DDH range	DDH best value	DZ	FCDH
$h_{\pi}^1$	$0 \rightarrow 30$	+12	+3	+7
$h_{\rho}^0$	$30 \rightarrow -81$	-30	-22	-10
$h_{\rho}^1$	$-1 \rightarrow 0$	-0.5	+1	-1
$h_{\rho}^2$	$-20 \rightarrow -29$	-25	-18	-18
$h_{\omega}^0$	$15 \rightarrow -27$	-5	-10	-13
$h_{\omega}^1$	$-5 \rightarrow -2$	-3	-6	-6

All values are quoted in units of  $g_{\pi} = 3.8 \times 10^{-8}$ .

DZ: Dubovik VM, Zenkin SV. *Ann. Phys.* 172:100 (1986)

FCDH: Feldman GB, Crawford GA, Dubach J, Holstein BR. *Phys. Rev. C* 43:863 (1991)

# DDH Model - Benchmark

In general, a measured PV NN observable can be expanded in terms of these:

$$O_{PV} = a_{\pi}^1 h_{\pi}^1 + a_{\rho}^0 h_{\rho}^0 + a_{\rho}^1 h_{\rho}^1 + a_{\rho}^2 h_{\rho}^2 + a_{\omega}^0 h_{\omega}^0 + a_{\omega}^1 h_{\omega}^1$$



E. G. Adelberger and W. C. Haxton, Ann. Rev. Nucl. Part. Sci. 35, 501 (1985).

DDH Weak Coupling	$(A_{\gamma}) np \rightarrow d\gamma$	$(A_{\gamma}) nd \rightarrow t\gamma$	$(\phi_{pV}) n-p$ ( $\mu\text{rad}/\text{m}$ )	$(\phi_{pV}) n-\alpha$ ( $\mu\text{rad}/\text{m}$ )	$(\frac{\Delta\sigma}{\sigma}) p-p$	$(\frac{\Delta\sigma}{\sigma}) p-\alpha$	$(A^p_Z) n^3\text{He} \rightarrow tp$
$a_{\pi}^1$	-0.107	-0.92	-3.12	-0.97	0	-0.340	-0.182
$a_{\rho}^0$	0	-0.50	-0.23	-0.32	0.079	0.140	-0.145
$a_{\rho}^1$	-0.001	0.103	0	0.11	0.079	0.047	0.0267
$a_{\rho}^2$	0	0.053	-0.25	0	0.032	0	0.0012
$a_{\omega}^0$	0	-0.160	-0.23	-0.22	-0.073	0.059	-0.1269
$a_{\omega}^1$	0.003	0.002	0	0.22	0.073	0.059	0.0495

Viviani *et al.*  
preliminary


*Experimental results generally agree with the DDH ranges, but:*

- *Uncertainties are large*
- *Some experimental results produce conflicting values for coupling constants (e.g. Values for  $h_\pi^1$  from  $^{18}\text{F}$  and  $^{133}\text{Cs}$  differ by several  $\sigma$ )*




p-p scat. 15, 45 MeV  $A_z^{pp}$   
 p-p scat. 221 MeV  $A_z^{pp}$   
 p- $\alpha$  scat. 46 MeV  $A_z^{pp}$

$^{133}\text{Cs}$ ,  $^{205}\text{Tl}$  anapole moments



$$\begin{array}{l}
 \mathbf{n+p \rightarrow d+\gamma} \quad \mathbf{A_\gamma^d} \\
 \mathbf{n-\alpha} \quad \mathbf{spin\ rot.} \quad \mathbf{d\phi^{n\alpha}/dz}
 \end{array}$$

*Unfortunately, the connection between the PV observables and the SM is essentially unknown.*



*with  $n^3\text{He}$   
(preliminary)*


$$\begin{array}{l}
 n+p \rightarrow d+\gamma \quad A_\gamma^d \\
 n-\alpha \quad \text{spin rot.} \quad d\phi^{n\alpha}/dz
 \end{array}$$

*Unfortunately, the connection between the PV observables and the SM is essentially unknown.*

# EFT Calculations (Upshot)

The  $\Delta S=0$  HWI can be parameterized in terms of 5 (8 with pions) low energy phenomenological constants.

At very low momenta ( $\leq \sim 50$  MeV) the constants essentially reduce to the 5 Danilov parameters:

  $\lambda_s^{0,1,2}, \lambda_t, \rho_t$

originally determined from NN scattering theory (Born approximation) write down simplest S-P amplitudes with PV and CP cons. amplitudes in addition to singlet and triplet strong ...

At higher momentum include explicit pions:  $h'_{\pi NN}, k_{\pi NN}^{1a}, \tilde{C}_{\pi}, \tilde{C}_{2\pi}$

# EFT Calculations

Write down 12 possible general P violating and CP conserving current-current terms with all isospin changes up to  $\Delta I=2$ :

$$O_1 = \frac{g_1}{\Lambda_\chi^2} \bar{\Psi}_N \mathbf{1} \gamma_\mu \Psi_N \bar{\Psi}_N \mathbf{1} \gamma^\mu \gamma_5 \Psi_N \quad O_2 = \frac{g_2}{\Lambda_\chi^2} \bar{\Psi}_N \mathbf{1} \gamma_\mu \Psi_N \bar{\Psi}_N \tau_3 \gamma^\mu \gamma_5 \Psi_N$$

$$\tilde{O}_1 = \frac{\tilde{g}_1}{\Lambda_\chi^3} \bar{\Psi}_N \mathbf{1} i \sigma_{\mu\nu} \mathbf{q}^\nu \gamma_\mu \Psi_N \bar{\Psi}_N \tau_3 \gamma^\mu \gamma_5 \Psi_N \quad \bullet \bullet \bullet \quad \text{etc...}$$

The NN contact potentials are expressed in terms of 12 parameters, but no mesons:

$$C_{1-5} = \frac{\Lambda_\chi}{2m_N} g_{1-5} \quad , \quad \tilde{C}_{1-5} = \tilde{g}_{1-5} + \frac{\Lambda_\chi}{2m_N} g_{1-5}$$

$$C_6 = \tilde{g}_6 - \frac{\Lambda_\chi}{2m_N} g_6$$

*Shi-Lin Zhu, et al. , Nuclear Physics A 748 (2005) 435-498*



Appropriate linear combinations of these produce the 5 Danilov coupling constants to be determined by experiment:

$$\begin{array}{ll}
 \lambda_7 \propto (C_1 - 3C_3) - (\tilde{C}_1 - 3\tilde{C}_3) & \\
 \lambda_s^0 \propto (C_1 + C_3) + (\tilde{C}_1 + \tilde{C}_3) & {}^3S_1 \rightarrow {}^1P_1 \quad I = 0 \\
 \lambda_s^1 \propto (C_2 + C_4) + (\tilde{C}_2 + \tilde{C}_4) & {}^1S_0 \rightarrow {}^3P_0 \quad I = 1 \\
 \lambda_s^2 \propto -\sqrt{\frac{8}{3}}(C_5 + \tilde{C}_5) & \\
 \rho_t \propto \frac{1}{2}(C_2 - C_4) + C_6 & {}^3S_1 \rightarrow {}^3P_1 \quad I = 1 \rightarrow 0
 \end{array}$$

$$\lambda_s^{pp} = \lambda_s^0 + \lambda_s^1 + \frac{1}{\sqrt{6}} \lambda_s^2$$



$$\lambda_s^{np} = \lambda_s^0 - \frac{2}{\sqrt{6}} \lambda_s^2$$

$$\lambda_s^{nn} = \lambda_s^0 - \lambda_s^1 + \frac{1}{\sqrt{6}} \lambda_s^2$$

# Experimental Program

We need at least 8 few body experiments to completely determine the EFT parameters.

Some have already been done:

*Longitudinal Asymmetries in p-p scattering:*

$$A_L^{pp}(13.6 \text{ MeV}) = -(0.93 \pm 0.20 \pm 0.05) \times 10^{-7} = -0.48 \lambda_s^{pp} m_N$$

Bonn: P.D. Evershiem *et al.* Phys. Lett. 256 (1991) 11

$$A_L^{pp}(45 \text{ MeV}) = -(1.5 \pm 0.22) \times 10^{-7} = -0.82 \lambda_s^{pp} m_N$$

PSI: S. Kistryn *et al.* Phys. Lett. 58 (1987) 1616  
R. Balzer *et al.* Phys. Rev. C. 30 (1984) 1409

# Experimental Program


## The TRIUMF 220 MeV pp experiment


$$A_L^{pp}(221 \text{ MeV}) = -(0.84 \pm 0.29 \pm 0.17) \times 10^{-7} \propto h_\omega^0 + h_\omega^1 \equiv h_\omega^{pp}$$

TRIUMF:

A.R. Berdoz *et al.* Phys. Rev. C 68 034004 (2003)

*Of order  $Q^3$  in EFT  
- no calculation yet ?*





**A.R. Berdoz *et al.* Phys. Rev. C 68 034004 (2003)**

# Experimental Program

## Longitudinal Asymmetry in $p$ - $\alpha$ scattering:

$$A_L^{p\alpha}(46 \text{ MeV}) = -(3.3 \pm 0.9) \times 10^{-7} = \left[ -0.48 \left( \lambda_s^{pp} + \frac{1}{2} \lambda_s^{np} \right) - 0.107 \left( \rho_t + \frac{1}{2} \lambda_t \right) \right] m_N$$

Bonn: J. Lang *et al.* Phys. Rev. Lett. 54 (1985) 170

## New experiments:

### ➤ Longitudinal asymmetry in proton scattering:

#### ▪ $p$ - $d$ :

$$A_L^{pd}(15 \text{ MeV}) = \left( -0.21 \rho_t - 0.07 \lambda_s^{pp} - 0.13 \lambda_t + 0.04 \lambda_s^{np} \right) m_N$$

# Experimental Program

*New experiments (or repeats):*

➤ *Neutron capture:*

▪ *Circ. Polarization:*

$$P_\gamma = (0.63\lambda_\gamma - 0.16\lambda_s^{np})m_N \quad \text{Very challenging!}$$

▪ *Gamma Asymmetry in np radiative capture:*

$$A_\gamma = -0.107 \rho_\gamma m_N \quad \text{LANSCE compl. SNS 2010}$$

▪ *Gamma Asymmetry in nd radiative capture:*

$$A_\gamma = (1.42\rho_\gamma + 0.59\lambda_s^{nn} + 1.18\lambda_\gamma + 0.51\lambda_s^{np})m_N \quad \text{Hard, SNS planned}$$

▪ *Proton Asymmetry in  $n^3\text{He}$  capture*

$$A_Z^p = (?)m_N \quad \text{Relatively easy, SNS approved ~2011}$$

# Experimental Program

*New experiments (or repeats):*

➤ *Neutron spin rotation:*

▪ *In helium:*

$$\frac{d\phi^{n\alpha}}{dz} = \left[ 1.2 \left( \lambda_s^{nn} + \frac{1}{2} \lambda_s^{np} \right) - 2.68 \left( \rho_t - \frac{1}{2} \lambda_t \right) \right] m_N \left[ \frac{\text{rad}}{m} \right]$$

W.M. Snow *et al*,  
Completed (NIST)

▪ *In hydrogen LH2*

$$\frac{d\phi^{np}}{dz} = \left[ 0.45 \lambda_s^{nn} + 1.28 \lambda_s^{np} + 0.45 \lambda_s^{pp} + 1.26 \rho_t - 0.63 \lambda_t \right] m_N \left[ \frac{\text{rad}}{m} \right]$$

SNS planned

# *Hadronic Parity Violation with Cold Neutrons*

*Two experiments (at the SNS):*

*NPDGamma:*

*Transversely polarized cold neutrons on hydrogen - looks for a directional asymmetry in the number of  $\gamma$ -rays, after decay:  $n + p \rightarrow d + \gamma$*

*n3He:*


*Longitudinally polarized cold neutrons on helium 3 - looks for a directional asymmetry in the number of protons after breakup:  $n + {}^3\text{He} \rightarrow t + p$*



# *Spallation Neutron Source (SNS)*




# *SNS Target Region*



# The Neutron Energy Scale


$$\lambda \approx \frac{29}{\sqrt{E_k}} \text{ fm}$$




$$E_k = \frac{\hbar^2 k^2}{2M_n} = k_B T \quad , \quad k = 43.4 \sqrt{E_k [\text{MeV}]} \text{ MeV} / c$$

# SNS Beam Properties


- total  $\sim 1.1 \times 10^{11}$  neutrons/second
- $4.1 \times 10^{10}$  n/s ,  $5.4 \times 10^{10}$  n/s,  $1.1 \times 10^{10}$  n/s for three example regions with no frame overlap
- 4 choppers required for various experimental conditions
  - eliminate overlap with slower neutrons from previous pulses
  - accommodate extraction of 0.89 nm beam
  - avoid potential background problems from leakage of fast neutrons
  - neutrons above 4.0 nm are not necessarily caught by this chopper arrangement (these come  $\sim 180$  ms after pulse onset (  $> 10$  frames later) intensity down by 4 orders of magnitude



# *SNS Ramp-Up Plan*



FNPB - 03/12/2009



Egal wo Du bist, es gibt immer ein...





Reciba estos chistes para adultos, gratis, email a [tavojaja@alta@elista.net](mailto:tavojaja@alta@elista.net)

Tavojaja  
Abr 2003

Dr. Tavo

# The Fundamental Neutron Physics Beam (FnPB)


- LH2 moderator
- 15 m long guide ~ 18 m to experiment
- one polyenergetic cold beam line
- one monoenergetic (0.89 nm) beam line
- ~ 40 m to nEDM UCN source
- 4 frame overlap choppers
- 60 Hz pulse repetition








# *Cold Beamline - Realized*




# *Flight path 13 - top view*



# The NPDGamma Collaboration

Los Alamos National Laboratory,  
**University of Manitoba,**  
University of Michigan,  
University of Tennessee,  
TJNAF,  
University of Dayton,  
Institute for Nuclear Research, Dubna,  
NIST,  
University of Kentucky



Indiana University,  
**TRIUMF,**  
University of New Hampshire,  
Oak Ridge National Laboratory,  
University of California-Berkeley,  
Hamilton College,  
KEK National Laboratory, Japan  
University of Virginia,  
UNAM



# The NPDGamma Observable / Theory

The main NPDGamma observable is the up-down asymmetry in the angular distribution of gamma rays with respect to the neutron spin direction.

$$\frac{d\sigma}{d\Omega} \propto \frac{1}{4\pi} (1 + A_\gamma \cos \theta)$$



$$A_{raw} = (P_n F_n D_n G) A_\gamma \cos \theta = \frac{1}{2} \left( \frac{\sigma_U^\uparrow - \sigma_D^\uparrow}{\sigma_U^\uparrow + \sigma_D^\uparrow} + \frac{\sigma_U^\downarrow - \sigma_D^\downarrow}{\sigma_U^\downarrow + \sigma_D^\downarrow} \right)$$

The observed cross-section is the result of an electro-magnetic transition between initial and final two nucleon states.

The possible amplitudes include both **parity even M1** and **parity odd E1 transitions** from L=1 states as a result of the weak perturbation.

$$\frac{d\sigma}{d\Omega} \propto \left| \langle \psi_f | \mathbf{E1} | \psi_i \rangle + \langle \psi_f | \mathbf{M1} | \psi_i \rangle \right|^2$$

$$H = H_s + V_{PNC} \quad a = \frac{\langle \psi_1 | V_{PNC} | \psi_0 \rangle}{\Delta E} \quad |\psi_{i,f}\rangle = |\psi_0\rangle + a|\psi_1\rangle$$

A measurement of the asymmetry at the 20 % level (10 ppb) will be the most precise measurement of the weak-pion nucleon coupling

$$\frac{ig_{\pi NN}h_{\pi}^1}{\sqrt{32M}} [\vec{\tau}_1 \times \vec{\tau}_2]_z [\vec{\sigma}_1 + \vec{\sigma}_2] \cdot \left[ \vec{p}, \frac{e^{-mr}}{4\pi r} \right]$$

$$\frac{g_{\pi NN}h_{\pi}^1}{\sqrt{32}} \approx 1.1 \times 10^{-6}$$

$$A_{\gamma} = -0.107 h_{\pi}^{\Delta I=1} \approx -0.107 \times 12 \times g_{\pi} = -5 \times 10^{-8}$$

# *NPDGamma EFT Relevance*

Systematic study of the NN weak interaction described in terms of a model independent theory appropriate at the low energy scale.

NN weak interaction effects enter into nucleon structure (needed for standard model tests) and atomic parity violation measurements.

5 EFT parameters :  $(\lambda_t, \lambda_s^{I=0,1,2}, \rho_t)$

Correspond to:  ${}^3S_1(I = 0) \leftrightarrow {}^1P_1(I = 0)$

${}^1S_0(I = 0, 1, 2) \leftrightarrow {}^3P_0(I = 0, 1, 2)$

${}^3S_1(I = 0) \leftrightarrow {}^3P_1(I = 1)$



NPDGamma asymmetry

relation to EFT constant:

$$A_{\gamma}^{\bar{n},p}(th.) = -0.107 m_N \rho_t = -5 \times 10^{-8}$$

*C.-P. Liu, Nuclear Physics Phys. Rev. C 75, 065501 (2007)*


# *LH<sub>2</sub> target and CsI detector array*



20L vessel of liquid parahydrogen

Ortho-hydrogen scatters the neutrons and leads to beam depolarization

- $3\pi$  acceptance
- Current-mode experiment
- $\gamma$ -rate  $\sim 100\text{MHz}$  (single detector)







**NPDGamma has successfully taken 48 days of continuous production data in 2006 - now on par with the best previous measurement - in preparation for one more year of production data at the SNS.**

# *Data Summary from 2006 run*

Number of good runs (8.5min long)	~5000
Neutron Polarization	$53 \pm 2.5\%$
Spin Flip Efficiency	$98.8 \pm 0.5\%$
Para fraction in LH <sub>2</sub> target	$99.98 \pm 0.2\%$
Al background	~25% (ave)
Depolarization	2%
Stern-Gerlach steering Asym	$10^{-10}$
$\gamma$ -ray circ.pol. Asym	$10^{-10}$

# Data Summary from 2006 run

Number of good runs (8.5min long)	~5000
Neutron Polarization	$53 \pm 2.5\%$
Spin Flip Efficiency	$98.8 \pm 0.5\%$
Para fraction in LH <sub>2</sub> target	$99.98 \pm 0.2\%$
Al background	~25% (ave)


Neutron polarization

2%





sym

$10^{-10}$

$10^{-10}$



# 2006 Hydrogen Results:




**Total statistical error:**

$$A_{\gamma,UD} = (-1.1 \pm 2.1) \times 10^{-7}$$

$$A_{\gamma,LR} = (-1.9 \pm 2.0) \times 10^{-7}$$

**Total systematic error: a (very) conservative 10% mostly due to pol.**

# Preliminary Hydrogen Results:



# *What's new for the SNS run*

- Supermirror Polarizer replaces the  $^3\text{He}$  Polarizer (x4.1)
- Higher moderator brightness (x12) => more cold/slow neutrons
- New LH2 target - thinner windows, smaller background contribution

Predicted size  $-5 \times 10^{-8}$  - NPDGamma will make a 20% measurement, most precise so far

- Installation begins in July 2009
- Production Hydrogen Data: Summer 2010

# *The Parity Violating Longitudinal Asymmetry in Polarized Cold Neutron Capture on Helium 3*

*$n^3\text{He}$*

**J.D. Bowman, S.I. Penttilä**

**R. Carlini**

**M. Gericke, S.A. Page**

**C. Crawford**

**V. Gudkov**

**J. Martin**

**C. Gillis**

**C. Gould**

**P-N. Seyo**

**P. Alacorn, T. Balascuta**

**S. Baessler**

**M. Viviani**

**Anna Hayes, Gerry Hale,  
and Andi Klein**

**Oak Ridge National Laboratory**

**Jefferson National Laboratory**

**University of Manitoba**

**University of Kentucky**

**University of South Carolina**

**University of Winnipeg**

**Indiana University**

**NC State University**

**Duke**

**Arizona State University**

**University of Virginia**

**INFN, Sezione di Pisa**

**Los Alamos National Laboratory**



# *$n^3\text{He}$ Principle of Measurement*

*Measure the asymmetry in the number of forward going protons in a  $^3\text{He}$  wire chamber as a function of neutron spin:*

$\vec{\sigma}_n \cdot \vec{k}_T$  *Directional PV asymmetry in the number of tritons*


$\vec{\sigma}_n \cdot \vec{k}_p$  *Directional PV asymmetry in the number of protons*  
*(much larger track length)*

- wire chamber is both target and detector*
- wires run vertical or horizontal*
- no crossed wire: keep the field simple to avoid electron multiplication (non-linearities)*





# Experimental Setup



- *longitudinal holding field - suppressed PC asymmetry*
- *RF spin flipper - negligible spin-dependent neutron velocity*
- *$^3\text{He}$  ion chamber - both target and detector*

- **MC simulations of sensitivity to proton asymmetry**




- including wire correlations

$$\delta A_{ph} = \frac{1}{\sqrt{NP_N}} \sqrt{\sigma_D^2 + \sigma_{coll}^2}$$

$$\sigma_d \simeq 6$$

- **tests at LANSCE FP12**

- fission chamber flux calibration
- prototype drift chamber R&D
- new beam monitors for SNS



- **MC simulations of sensitivity to proton asymmetry**




- including wire correlations

$$\delta A_{ph} = \frac{1}{\sqrt{NP_N}} \sqrt{\sigma_D^2 + \sigma_{coll}^2}$$





$$\sigma_d \simeq 6$$

- **tests at LANSCE FP12**


- fission chamber flux calibration
- prototype drift chamber R&D
- new beam monitors for SNS



# LANSCCE $n^3\text{He}$ Chamber Tests










*Right: LANL FP12 beam line with new beam monitor installed*

*Below: New beam monitor signal at 100  $\mu$ A proton beam current.*



# $n^3\text{He}$ Relevance

Systematic study of the NN weak interaction described in terms of a model independent theory appropriate at the low energy scale.

NN weak interaction effects enter into nucleon structure (needed for standard model tests) and atomic parity violation measurements.

5 EFT parameters :  $(\lambda_t, \lambda_s^{I=0,1,2}, \rho_t)$

Correspond to:

$${}^3S_1(I = 0) \leftrightarrow {}^1P_1(I = 0)$$

$${}^1S_0(I = 0, 1, 2) \leftrightarrow {}^3P_0(I = 0, 1, 2)$$

$${}^3S_1(I = 0) \leftrightarrow {}^3P_1(I = 1)$$

$n^3\text{He}$  asymmetry  
relation to EFT constant:

$$A_p^{\bar{n}, {}^3\text{He}}(th.) = \kappa \lambda_s^{I=0} \approx 3 \times 10^{-7}$$

*M. Viviani, R. Schiavilla, calculation in progress*



# Status/Schedule

- *n-<sup>3</sup>He experiment approved by the FnPB PRAC, 2008-01-07*
  - *first measurement of PV in the n-<sup>3</sup>He reaction*
  - *large asymmetry ~10<sup>-7</sup>*
  - *proposed measurement accuracy*  $\delta A = 1.0 \times 10^{-8}$
- *recent progress in experimental design*
  - *full 4-body calculation of PV observable*
  - *R&D projects on target/detector design at LANL*
  - *new spin flipper design permitting compact / less expensive layout*
  - *preliminary holding field design*
- *leverage existing hardware / technology*
  - *major components based on similar NPDGamma instrumentation*
  - *can reuse NPDGamma electronics / power supplies*
- *FnPB infrastructure*
  - *no safety hazards, no LH<sub>2</sub> target, new power or cooling requirements*
  - *minimal modification of FnPB cave - stand for n-<sup>3</sup>He solenoid*
  - *technician support for readiness review preparation, setup of experiment*
- *Tentative Schedule*
  - *2009-2010 Design and development*
  - *Late 2011 Installation*
  - *2012 Run*

*Ciao*